arXiv:1607.07055v1 [math.GR] 24 Jul 2016

CAYLEY PARAMETRIZATION AND THE ROTATION GROUP OVER A NON-ARCHIMEDEAN PYTHAGOREAN FIELD

M. G. MAHMOUDI

Abstract. Using Cayley transform, we show how to construct rotation matrices *infinitely near* the identity matrix over a non-archimedean pythagorean field. As an application, an alternative way to construct non-central proper normal subgroups of the rotation group over such fields is provided.

Mathematics Subject Classification: 11E57, 12D15, 20G15, 65F35.

Keywords: Cayley parametrization; Frobenius norm; orthogonal group; rotation group; pythagorean field; non-archimedean field.

1. Introduction

The Cayley transform provides an important parametrization of the rotation group of the euclidean space. This map which was introduced by Arthur Cayley in 1846, associates a rotation matrix to every skew symmetric matrix of eigenvalues different from -1. The Cayley parameterization was used by J. Dieudonné in [3, §15, §16] to construct normal subgroups of the rotation group of a vector space V of dimension $n = 3$ equipped with an anisotropic quadratic form over the field of p-adic numbers or the field of formal Laurent series in one variable over the field of real numbers for arbitrary $n \geqslant 3$. Later, using the properties of *Elliptic spaces*, E. Artin in [1, Ch. V, §3] constructed non-central subgroups of the rotation group $\mathrm{SO}(V)$ of a vector space V of dimension $n \geqslant 3$ equipped with an anisotropic quadratic q form over a field F with a non-archimedean ordering.

We simplify the Artin construction using the Cayley parametrization. The main new observations that make this simplification possible are the following:

(1) Assume that n is odd or n is even and $\det q$ is trivial. If F is not formally real pythagorean then the rotation group $\mathrm{SO}(V)$ is never projectively simple (see (3.1) and (3.2)). This enables us to concentrate on the case where the base field F is pythagorean, hence the Frobenius norm is at disposal and one may consider it instead of the maximum norm as used in [1, Ch. V, §3].

(2) If F is a formally real pythagorean field with a non-archimedean ordering then one can find a rotation matrix $A \in M_n(F)$ such that the Frobenius norm of $A - I \neq 0$ is infinitely small (see (3.3)).

Finally, as another application, we complement a result due to L. Bröcker [2], which implies that the orthogonal groups $\mathrm{SO}(V)$ are all projectively simple for the case where the Witt index ν of q is zero, $n \geqslant 3$ and $n \neq 4$ precisely in the case where F is pythagorean formally real field, admits only of archimedean ordering and q is similar to $x_1^2 + x_2^2 + \cdots + x_n^2$. This addresses the following passage by J. Dieudonné in [3, p. 39]: *"... la simplicité du groupe des rotations (pour $\nu = 0$, $n > 2$ et $n \neq 4$) sur le corps des nombres réels $\mathbb{R}$, apparaît comme un phénomène très particulier, dû à la structure très spéciale du corps $\mathbb{R}$ parmi les corps commutatifs. Il y aurait lieu de rechercher s'il existe d'autres corps qui partagent avec lui cette propriété."*.

2. Notation and Terminology

Throughout this paper, F denotes a field of characteristic different from two. Let (V, b) be bilinear space of dimension n over a field F and let $q : V \to F$ be its associated nondegenerate quadratic form given by $q(x) = b(x, x)$. The determinant of b is the determinant of a Gram matrix of b modulo $F^{\times 2}$ and is denoted by $\det b$ or $\det q$. One says that b or q represents a scalar $\lambda \in F$ if there exists a nonzero vector $v \in V$ such that $b(v, v) = \lambda$. A vector $u \in V$ is called *anisotropic* (resp. *isotropic*) if $q(u) \neq 0$ (resp. $q(u) = 0$). A quadratic form q is said to be *anisotropic* if all nonzero vectors in V are anisotropic, otherwise q is called *isotropic*. The *Witt index* ν of (V, b) is defined as the maximal dimension of a totally isotropic subspace of V. As the characteristic of F is different from 2, there exists an orthogonal basis $\{e_1, \cdots, e_n\}$ of (V, b). It is known that for $n \geqslant 3$ the group $\mathrm{SO}(V)$ is a non-abelian group and its center is either $\{\pm \mathrm{id}\}$ or $\{\mathrm{id}\}$ depending on the parity of n (see [4, p. 51]).

For every $\sigma \in \mathrm{O}(V)$ by the Cartan-Dieudonné theorem, there exists a decomposition $\sigma = \prod_{i=1}^{m} \tau_{u_i}$ where $m \leqslant n$ and τ_{u_i} is a reflection along the anisotropic vectors $u_i \in V$ given by

$$\tau_{u_i}(x) = x - 2\frac{b(x, u_i)}{q(u_i)}u_i.$$

It is known that the class of $\theta(\sigma) = \prod_{i=1}^{n} q(u_i)$ in the quotient group $F^{\times}/F^{\times 2}$, which is called the *spinor norm* of σ, is independent of the choices and the number of u_i's.

We recall that a field F is said to be *formally real* (or *ordered field*) if -1 is not a sum of squares in F. A field F is called *pythagorean* if every sums of squares is again a square. An element c of a formally real field F is called *totally positive* if c is positive with respect to all orderings of F.

If F is a pythagorean field then for every matrix A with entries in F, the *Frobenius norm* of A, denoted by $\|A\|$, is defined as the absolute value of the square root of the sum of squares of entries of A. The Frobenius norm satisfies the inequalities $\|AB\| \leqslant \|A\|\|B\|$ and $\|A + B\| \leqslant \|A\| + \|B\|$ when A and B are of appropriate sizes.

For any matrix $A \in M_n(F)$ such that $I + A$ is invertible, the map $A \mapsto (I - A)(I + A)^{-1}$ is called the *Cayley map*. For any skew-symmetric matrix $A \in M_n(F)$ such that the matrix $I+A$ is invertible, the matrix $Q = C(A)$ satisfies $Q^TQ = I$ and $\det Q = 1$. Conversely for every orthogonal matrix Q such that $I + Q$ is invertible $C(Q)$ is a skew-symmetric matrix, see [6, p. 56].

3. Rotation group over non-archimedean fields

As a first observation we have the following result:

Proposition 3.1. *Let (V, b) be anisotropic bilinear space of dimension $n \geqslant 2$ over a field F. Then the image of the spinor norm $\theta : \mathrm{SO}(V) \to F^{\times}/F^{\times 2}$ is trivial if and only if F is a formally real pythagorean field and q represents only one square class in $F^{\times}$, in particular q is similar to the quadratic form $x_1^2 + \cdots + x_n^2$.*

Proof. First assume that the spinor norm map is the trivial map. Let $u \in V$ be an anisotropic vector with $q(u) = d \neq 0$. Then for any anisotropic vector $v \in V$ as $\theta(\tau_u\tau_v)$ is trivial, the scalars $q(u)$ and $q(v)$ are in the same square class in $F^{\times}$. It follows that q is isomorphic to $dx_1^2 + \cdots + dx_n^2$. Let $\alpha, \beta \in F$, we should prove that there exists $\gamma \in F$ such that $\gamma^2 = \alpha^2 + \beta^2$. As $n \geqslant 2$, we may consider two orthogonal vectors $u, v \in V$ with $q(u) = q(v) = d \neq 0$. Now consider the vector $w = \alpha u + \beta v \in V$. We have $q(w) = (\alpha^2 + \beta^2)d$. As $q(w)$ and $q(u)$ are in the same square classes, the quantity $\alpha^2 + \beta^2$ is a square in F. If F is not formally real,

then -1 is a sum of squares in F, hence is a square since F is pythagorean. As $q \simeq dx_1^2 + \cdots + dx_n^2$ and $n \geqslant 2$, the form q would be isotropic, contradiction. The converse follows from the Cartan-Dieudonné theorem and the fact that the current hypotheses imply that the spinor norm of the product of two arbitrary reflections is trivial. □

Proposition 3.2. *Let (V, b) be an anisotropic bilinear space of dimension $n \geqslant 3$ over a field F such that $\mathrm{SO}(V)$ is projectively simple. Then the spinor norm map $\theta : \mathrm{SO}(V) \to F^\times/F^{\times 2}$ is the trivial map precisely in the following cases (i) n is odd, (ii) n is even and the determinant of b is trivial.*

Proof. First suppose that θ is the trivial map. Let $\{e_1, \cdots, e_n\}$ be an orthogonal basis of (V, b). We have $-\mathrm{id} = \tau_{e_1}\tau_{e_2}\cdots\tau_{e_n}$. It follows that in the case where n is even, $-\mathrm{id} \in \mathrm{SO}(V)$ and $\theta(-\mathrm{id})$ which coincides with the determinant of b, is trivial.

Conversely suppose that (i) or (ii) hold. In the case (i) the group $\mathrm{SO}(V)$ has trivial center. Thus the hypotheses actually say that $\mathrm{SO}(V)$ is itself a simple group. It follows that $\ker(\theta)$ is either the trivial group or is the whole $\mathrm{SO}(V)$. The first case is impossible as $\mathrm{SO}(V)$ is non-abelian. The second case implies that $\mathrm{im}(\theta)$ is trivial. In the case (ii) the center of $\mathrm{SO}(V)$ is the subgroup $\{\pm\mathrm{id}\}$. As $-\mathrm{id} = \tau_{e_1}\tau_{e_2}\cdots\tau_{e_n}$ and the determinant of b is assumed to be trivial, $\theta(-\mathrm{id})$ is trivial as well. Therefore, the kernel of the spinor norm map θ is a subgroup of $\mathrm{SO}(V)$ containing $\{\pm\mathrm{id}\}$. As $\mathrm{SO}(V)$ is assumed to be projectively simple, $\ker(\theta)$ is either $\{\pm\mathrm{id}\}$, or $\mathrm{SO}(V)$. If $\ker(\theta) = \{\pm\mathrm{id}\}$, we have $\mathrm{SO}(V)/\{\pm\mathrm{id}\} \simeq \mathrm{im}(\theta)$. This shows that $\mathrm{SO}(V)/\{\pm\mathrm{id}\}$ is an abelian simple group. It follows that this quotient group is either the trivial group (thus $\mathrm{im}(\theta)$ is the trivial group) or a group of prime order. The later case is ruled out as a non-abelian group is never projectively cyclic. The former case also implies the triviality of $\mathrm{im}(\theta)$. □

Proposition 3.3. *Let F be a pythagorean field with a non-archimedean ordering $\leqslant$. Then there exists a matrix $A \in M_n(F)$ with $A \neq \pm I$, $A^T A = I$ and $\det A = 1$ such that $\|I - A\|$ is infinitely small.*

Proof. Let ϵ be an infinitely small positive element of F and let $B \in M_n(F)$ be a nonzero skew-symmetric matrix whose entries are rational. Consider the matrix $A = C(\epsilon B)$, where C is the Cayley map. We claim that $\|I - A\|$ is infinitely small. First note that $I + \epsilon B$ is invertible, hence $C(\epsilon B)$ is meaningful and is different from $\pm I$. Let m be an odd positive integer. The relation $I + \epsilon^m B^m = (I + \epsilon B)D$ where $D = (I - \epsilon B + \epsilon^2 B^2 + \cdots + \epsilon^{m-1} B^{m-1})$ implies that $\|(I + \epsilon B)^{-1} - D\| = \epsilon^m \|B^m (I + \epsilon B)^{-1}\|$. By the Cramer rule every entry of $B^m(I + \epsilon B)^{-1}$ is of the form $p(\epsilon)/q(\epsilon)$ where $p(X), q(X) \in \mathbb{Q}[X]$ are of degree at most n. It follows that for sufficiently large m, the quantity $\|(I + \epsilon B)^{-1} - D\| = \epsilon^m p(\epsilon)/q(\epsilon)$ is infinitely small. Hence

$$\begin{aligned}\|I - A\| =& \|I - (I - \epsilon B)(I + \epsilon B)^{-1}\| \\ \leqslant& \|I - (I - \epsilon B)D\| + \|(I - \epsilon B)((I + \epsilon B)^{-1} - D)\| \\ \leqslant& \|I - (I - \epsilon B)D\| + \|I - \epsilon B\|\|(I + \epsilon B)^{-1} - D\|\end{aligned}$$

Both quantities $\|(I - (I - \epsilon B))D\|$ and $\|(I + \epsilon B)^{-1} - D\|$ are infinitely small and the proof is complete. □

Proposition 3.4. *Let F be a pythagorean ordered field and let (V, b) be a bilinear space of dimension $n \geqslant 3$, isometric to $x_1^2 + \cdots + x_n^2$. If F carries a non-archimedean ordering then $\mathrm{SO}(V)$ contains a proper non-central normal subgroup.*

Proof. We claim that the group

$$N = \{\sigma \in \mathrm{SO}(V) : \ \|x - \sigma(x)\| \text{ is infinitely small if } q(x) = 1\}$$

is a non-trivial normal subgroup of $\mathrm{SO}(V)$ (the idea of considering this subgroup was borrowed from [5, p. 150]). To prove this claim we first show that N is a subgroup of $\mathrm{SO}(V)$. Consider two elements $\sigma, \tau \in N$. We should prove that $\sigma\tau$ also belongs to N. We have $\|x - \sigma\tau(x)\| = \|(x - \tau(x)) + (\tau(x) - \sigma\tau(x))\| \leqslant \|x - \tau(x)\| + \|\tau(x) - \sigma\tau(x)\| = \|x - \tau(x)\| + \|x - \sigma(x)\|$, which is infinitely small. The fact that $\sigma^{-1} \in N$ when $\sigma \in N$ and the normality of N is straightforward. It remains to show that N is a nontrivial subgroup of $\mathrm{SO}(V)$. Consider two orthogonal vectors $u, v \in V$ with $q(u) = q(v) = 1$. Let $\sigma = -\tau_u \in \mathrm{SO}(V)$. We have $\|v - \sigma(v)\| = \|v + \tau_u(v)\| = \|2v\| = 2$. Hence $\sigma \notin N$, thus $N \neq \mathrm{SO}(V)$. By (3.3), there exists a rotation matrix A such that $\|I - A\|$ is infinitely small. It follows that for every x with $\|x\| = 1$ we have $\|x - Ax\| \leqslant \|I - A\|$ is infinitely small. Hence $A \neq \pm I$ is an element of N and $N \neq \{\pm I\}$ □

Corollary 3.5. *(Bröcker) Let (V, b) be a bilinear space of dimension $n \geqslant 3$ over a field F whose Witt index is zero. Then the rotation group $\mathrm{SO}(V)$ is a projectively simple if and only if F is a pythagorean formally real field and q is similar to the quadratic form $x_1^2 + x_2^2 + \cdots + x_n^2$.*

Proof. In [2], Bröcker proves that if F is a pythagorean formally real field which admits only of archimedean orderings, $n \geqslant 3$ and $n \neq 4$ and q is anisotropic then the kernel of the spinor norm $\theta : \mathrm{SO}(V) \to F^\times/F^{\times 2}$ coincides with the commutator subgroup $\Omega(V)$ of the orthogonal group $\mathrm{O}(V)$ (see [2, (1.7)]) and the group $\Omega(V)$ is projectively simple (see [2, Satz (1.10)]). Of course when q represents only one square class, the whole rotation group $\mathrm{SO}(V)$ coincides with the kernel of the spinor norm. Hence the sufficiency of the conditions can be obtained from Bröcker's theorem. The sufficiency follows from (3.1), (3.2) and (3.4). □

Acknowledgments. The support from Sharif University of Technology is gratefully acknowledged.

References

[1] Artin, E. *Geometric algebra.* Interscience tracts in pure and applied mathematics. No. 3. New York: Interscience Publishers, Inc.; London: Interscience Publishers Ltd. (1957).

[2] Bröcker, L. Zur orthogonalen Geometrie über pythagoreischen Körpern. J. Reine Angew. Math. **268/269**, 68-77 (1974).

[3] Dieudonné, J. *Sur les groupes classiques.* Publications de l'Institut de Mathématique de l'Université de Strasbourg, VI. Actualités Scientifiques et Industrielles, No. 1040. Hermann, Paris, 1973.

[4] Dieudonné, J. *La géométrie des groupes classiques.* Ergebnisse der Mathematik und ihrer Grenzgebiete. Band 5. Berlin-Heidelberg-New York: Springer-Verlag (1971).

[5] Perrin, D. *Cours d'algèbre*, Paris: Ellipes (1996).

[6] Weyl, H. *The classical groups, their invariants and representations.* Reprint of the second edition (1946) of the 1939 original. Princeton University Press, Princeton, N.J. (1939).

M. G. Mahmoudi, mmahmoudi@sharif.ir,
Department of Mathematical Sciences, Sharif University of Technology, P. O. Box 11155-9415, Tehran, Iran.